Updates on Frozen Embryo Transfer



Carrie E. Bedient, MD, FACOG

Clinical Associate Professor Kirk Kerkorian School of Medicine Director of Reproductive Endocrinology & Infertility Mountainview Ob/Gyn Residency University of Nevada, Las Vegas Fertility Center of Las Vegas Las Vegas, NV

Practices surrounding IVF cycles have evolved widely over the past 40 years. Cutting edge techniques several years ago have naturally been replaced by different protocols and procedures to take advantage of improved safety and efficacy profiles. Through application of vitrification techniques over slow freezing, embryo viability post thaw is significantly improved, creating space for other advantages of frozen embryo transfers (FET) to emerge. FETs improve success rates, perinatal risk profiles and maternal risk profiles. The advent of frozen embryo transfers collectively improved success rates as well as decreased pregnancy loss, both ultimately improving number of live births.

Factors contributing to this increased success vary. One beneficial aspect of freeze-all cycles includes the ability to push oocyte development longer, rather than triggering when the first few follicles become large. This allows for more aggressive stimulation with higher oocyte yields. Studies have shown that while 19mm is an optimal size for retrieval, larger follicles yield good quality blasts in similar percentages. With a frozen transfer physicians can stimulate the ovaries more aggressively, particularly in high responder patients, without risking severe OHSS due to high numbers of follicles, higher estrogen levels, and the threat of continued elevated HCG in the setting of an ongoing pregnancy. Freezeall cycles also permit preimplantation genetic testing (PGT) to be performed, and embryo selection for transfer improves with potential combination of multiple cohorts of available embryos. PGT tested embryos ensure obviously aneuploid embryos are not transferred. The impact of transferring clearly aneuploid embryos affects not only patient distress in an adverse outcome, but also the potential resultant desire for termination of an abnormal pregnancy that could have been avoided. Genetic testing also permits greater comfort in transferring a single embryo, thereby cutting risk of multiple pregnancies down to spontaneous levels instead of the historically elevated multiple gestations associated with IVF.

Optimization of embryo-endometrial synchrony likely greatly contributes to improved success rates. With the presence of ovarian stimulation immediately prior to transfer, the endometrial development is altered, negatively impacting timing of appropriate endometrial histology and hormonal receptor regulation. The relative risks of perinatal complications significantly decrease with FETs. This is partly attributable to improved success rates and ability to do single embryo transfer of euploid embryos which minimizes the incidence of multiple gestations. In addition to the increase of singleton pregnancies, avoidance of perinatal complications remains significant even when comparing outcomes in singleton fresh vs. frozen cycles. Placental diseases, low birthweight and prematurity, and ovarian hyperstimulation syndrome all decrease with frozen embryo cycles. That said, hypertensive disorders of pregnancy are definitively higher with FETs and that factor should be accounted for while counseling patients. Cumulatively, FET increases success rates and safety profiles for patients due to increased ability to optimize oocyte stimulation practices, embryonic genetic testing, endometrial development, and timing of embryo transfer.